



SURVEILLANCE UPDATE ON NEONATAL TETANUS
Morbidity Week 11 (January 1 - March 15), Philippines 2015



| REGION | Confirmed NT Case | *Mothers with ≥ TT2 | Place of Delivery | | | Cord Cutting Tool Used | | | | Delivery Attendants | | | | | Outcome | |
|------------|-------------------|---------------------|-------------------|----------------|----------|------------------------|----------------|----------------|----------|---------------------|----------------|----------------|------------------------------|----------|----------------|----------------|
| | | | Hospital/Clinic | Home | Others | Scissor | Blade | Bamboo | Others | Physician | Nurse | Midwife | Traditional Birth Attendants | Others | Died | Alive |
| I | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| II | 1 | 1 (100%) | 0 | 1 (100%) | 0 | 1 (100%) | 0 | 0 | 0 | 0 | 0 | 0 | 1 (100%) | 0 | 1 (100%) | 0 |
| III | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| IVA | 1 | 0 | 0 | 1 (100%) | 0 | 0 | 0 | 1 (100%) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 (100%) |
| IVB | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| V | 3 | 1 (33%) | 1 (33%) | 2 (67%) | 0 | 1 (33%) | 0 | 2 (67%) | 0 | 0 | 1 (33%) | 0 | 2 (67%) | 0 | 1 (33%) | 2 (67%) |
| VI | 2 | 0 | 0 | 2 (100%) | 0 | 1 (50%) | 0 | 1 (50%) | 0 | 0 | 0 | 0 | 1 (100%) | 0 | 1 (100%) | 1 (100%) |
| VII | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| VIII | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| IX | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| X | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| XI | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| XII | 1 | 0 | 0 | 1 (100%) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 (100%) | 0 | 1 (100%) | 0 |
| ARMM | 1 | 0 | 0 | 1 (100%) | 0 | 0 | 1 (100%) | 0 | 0 | 0 | 0 | 0 | 1 (100%) | 0 | 0 | 1 (100%) |
| CAR | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| CRG | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| NCR | 1 | 0 | 0 | 1 (100%) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 (100%) | 0 | 0 | 1 (100%) | 0 |
| PHL | 10 | 2 (20%) | 1 (10%) | 9 (90%) | 0 | 3 (30%) | 1 (10%) | 4 (40%) | 0 | 0 | 1 (10%) | 1 (10%) | 6 (60%) | 0 | 5 (50%) | 5 (50%) |


*Mothers of confirmed NT cases

NOTE: Totals per variable (Place of Delivery, Cord Cutting Tool Used, Delivery Attendants, Outcome) may be inconsistent due to missing information.

Prepared by:


Lee Anne C. Mappatlo - Esguerra, RN
 National NT Surveillance Officer

Approved by:


Vito G. Roque Jr., RM, MD, PHSAE
 Chief, Public Health Surveillance & Informatics Division